

Anaphylaxis Emergency Action Plan

Patient Name:		Age:
Allergies:		
Asthma Yes (high risk for sever	e reaction) 🔲 No	
Additional health problems besides	anaphylaxis:	
Concurrent medications:		
MOUTH THROAT* SKIN GUT LUNG* HEART*	Symptoms of Anaphylaxis itching, swelling of lips and/or tonguitching, tightness/closure, hoarsene itching, hives, redness, swelling vomiting, diarrhea, cramps shortness of breath, cough, wheeze weak pulse, dizziness, passing out	ss
Only a few symptoms *Some s	may be present. Severity of symptoms symptoms can be life-threatening. ACT	can change quickly. FAST!
Emergency Action Steps - D 1. Inject epinephrine in thigh using (c	O NOT HESITATE TO GIVE EPINEPHRI heck one):	NE! Adrenaclick (0.3 mg)
·*)	☐ Auvi-Q (0.15 mg)	Auvi-Q (0.3 mg)
	EpiPen Jr (0.15 mg)	☐ EpiPen (0.3 mg)
	Epinephrine Injection, US ☐ (0.15 mg)	SP Auto-injector- authorized generi
	Other (0.15 mg)	Other (0.3 mg)
Specify others:		
IMPORTANT: ASTHMA INHALERS	AND/OR ANTIHISTAMINES CAN'T BE D	DEPENDED ON IN ANAPHYLAXIS.
2. Call 911 or rescue squad (before	calling contact)	*
3. Emergency contact #1: home	work	cell
Emergency contact #2: home	work	cell
Emergency contact #3: home	work	cell
Doctor's Signature/Date/Phone Numb	per	_
Parent's Signature (for individuals un	ider age 18 yrs)/Date	

This information is for general purposes and is not intended to replace the advice of a qualified health professional. For more information, visit www.aaaai.org. © 2013 American Academy of Allergy, Asthma & Immunology 7/2013